
Subject: Re: row calculation in a 2D array
Posted by [Martin Schultz](#) on Tue, 25 Aug 1998 07:00:00 GMT
[View Forum Message](#) <> [Reply to Message](#)

Jonas wrote:

>
> thanxalot (both Kevin and David), it helped. I wasn't aware of the function
> "TOTAL", still being a newbie and all.
>
> This was kind of a special example since total could do all the work. What
> if I would like to do another calculation where total is not applicable...
> in other words:
> is there generally an easy way to perform the same calculation over all the
> (sub-)rows in a 2D array?
>
> I do not have such a problem at the moment, but I assume it is likely that I
> will encounter it in the future (at least if i don't ask now :-)).
>
> sincerely
> Jonas Svensson

Well, if you want to compute averages first over rows 0:3, then 4:7,
then 8:11, etc. you may not get around using a loop. Although, in some
cases you could try your luck with `reform()` and a subsequent call to
Kevin's average routine (which I like very much: THANKS!). But this is
certainly prone to errors...

Martin.

--

Dr. Martin Schultz
Department for Earth&Planetary Sciences, Harvard University
109 Pierce Hall, 29 Oxford St., Cambridge, MA-02138, USA

phone: (617)-496-8318
fax : (617)-495-4551

e-mail: mgs@io.harvard.edu
Internet-homepage: <http://www-as.harvard.edu/people/staff/mgs/>
