Subject: Re: Faster than this...
Posted by Ricardo Fonseca on Fri, 04 Feb 2000 08:00:00 GMT
View Forum Message <> Reply to Message

Nope, not a trick question, just a vary basic one

Thanks, Ricardo

```
> From: Craig Markwardt < craigmnet@cow.physics.wisc.edu>
> Organization: U. Wisc. Madison Physics -- Compact Objects
> Reply-To: craigmnet@cow.physics.wisc.edu
> Newsgroups: comp.lang.idl-pvwave
> Date: 03 Feb 2000 19:00:34 -0600
> Subject: Re: Faster than this...
>
 Ricardo Fonseca <zamb@physics.ucla.edu> writes:
   I have this line of code in IDL 5.2a
>>
>> for j=0, N-1 do Data[j]=XAxisData[j]^2*Data[j]
>>
>> Where both Data and XAxisData have the same dimension N.
>> Does anyone know if there is a faster way of doing this
>> (i.e. avoiding the for cicle)?
 I hope this isn't a trick question. The expression
>
  Data = XAxisDAta^2 * Data
> should work fine. Don't worry about Data being overwritten. The
> right hand side is fully evaluated with the "old" value of Data,
 before the "new" value of Data is assigned.
>
> Craig
> Craig B. Markwardt, Ph.D. EMAIL: craigmnet@cow.physics.wisc.edu
> Astrophysics, IDL, Finance, Derivatives | Remove "net" for better response
```