Subject: Re: MPFIT/MPFIT2DFUN (Markwardt) Posted by Craig Markwardt on Sat, 23 Dec 2000 21:07:13 GMT View Forum Message <> Reply to Message

Greetings!

The function MPFIT2DFUN is designed to fit gridded 2-dimensional data. ie an image.

However, the task you are asking, a scalar function of two variables, is not very hard to do with the standard MPFITFUN. I even cover it as a frequently asked question on my web page (hint, hint...).

What you want to do is make a *single*, *two*-dimensional independent variable consisting of both your X and Y data. MPFITFUN doesn't care that your independent variable is of different dimension than the dependent variable. Your new independent variable would be constructed like this:

```
XY = [[X],[Y]]
```

You would call MPFITFUN with XY as your independent variable, and Z as your dependent variable. You need to make sure that your user function can split this independent data array apart:

```
function myfunc, xy, p
 x = xy(*,0)
 y = xy(*,1)
 ... continue with calculations ...
end
```

Good luck, Craig

endobeck@my-deja.com writes:

- > I have one-dimensional x, y, and z arrays, with x & y being positions
- > (independent variables), and z being an array of dependent variable
- > values. All 3 arrays have the same number of elements.
- > I'd like to fit a function of two variables (a plane) to the
- > data, and I've been trying to use MPFIT2DFUN. It seems I have to make
- > the x & y arrays into 2-d arrays, but I am not clear as to why or how.
- > Can someone fill me in?
- > > Cheers,
- > Eli

> Sent via Deja.com > http://www.deja.com/	
Craig B. Markwardt, Ph.D. EMAIL: craigmnet@cow.physics.wisc.edu	
Astrophysics, IDL, Finance, Derivatives Remove "net" for better response	