
Subject: Re: Hankel (Fourier-Bessel) Transform
Posted by [thompson](#) on Tue, 24 Jul 2001 19:15:17 GMT
[View Forum Message](#) <> [Reply to Message](#)

Georg.Pabst@nrc.ca (Georg Pabst) writes:

> Hi,

> I'm looking for the Hankel (Fourier-Bessel) Transform, i.e.,
> $\int_0^{\infty} f(t) \text{BesselJ}(t \cdot r) t \, dt$ being implemented in IDL.

> There is a paper "Siegman A. 1980. Quasi fast Hankel transform. Opt.
> Lett. 1, 13-15" and one can also find the code in Fortran or C...

> Thanks,
> Georg

Here's an old program that I think might be what you need.

Bill Thompson

```
FUNCTION HANKEL,F
;
; This function returns the Hankel transform of the argument.
;
S = SIZE(F)
IF S(0) NE 1 THEN BEGIN
  PRINT,'*** Variable must be a one-dimensional array, name= F, routine HANKEL.'
  RETURN,F
ENDIF
;
X = INDGEN(F)
K = ( 2. * !PI / FLOAT(N_ELEMENTS(X)) ) * X
SC = 0.*X + 1.
IF N_ELEMENTS(SC) GT 3 THEN BEGIN
  SC(0) = 3.D0 / 8.D0
  SC(1) = 7.D0 / 6.D0
  SC(2) = 23.D0 / 24.D0
ENDIF
;
H = BES0( K # X ) # ( K * F * SC )
;
RETURN,H
END
```
