
Subject: Re: Band reader

Posted by [julio](#) on Tue, 30 Dec 2003 13:13:15 GMT

[View Forum Message](#) <> [Reply to Message](#)

Thanks Fotis, I'll try it!

Julio

Fotis <fotis@physics.uoc.gr> wrote in message news:<[bsr9pk\\$njg\\$1@nic.grnet.gr](mailto:bsr9pknjg1@nic.grnet.gr)>...

> Julio wrote:

>> Hello, I'm a new user of ENVI/IDL and I'm writing a routine to
>> calibrate the channels 1 and 2 of AVHRR images. According to KLM User's
>> Guide, the slope and intercept coefficients values depend on the CN.
>> I'm using the ENVI_GET_IMAGE to read the channels, but it reads the
>> matrix, and not unit values. How can I do to read only the CN and then
>> apply the right values of intercept and slope?
>> Thanks a lot!

> Assume that calibr. factors change for CN values larger or less
> than 499.
> Assume b1 is the matrix of the first band (0.63 micron)
> temp1=fltarr(ns,nl)
> temp2=fltarr(ns,nl)
> calibr_b1=fltarr(ns,nl)
>
> temp1=float(b1)* SLOPE1 + INTERC1
> temp2=float(b1)* SLOPE2 + INTERC2
>
> calibr_b1=temp1 * (b1 le 499) + temp2 * (b1 gt 499)
