Subject: Re: combinatory analysis with restrictions Posted by Craig Markwardt on Thu, 09 Dec 2004 06:41:01 GMT View Forum Message <> Reply to Message

fleduc@lycos.com (Francois) writes:

```
> Hello,
```

- > Considering 3 variables (a,b,c) that can take 11 discrete values
- > between 0 and 1: [0, 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9,
- > 1.0]

Why don't you make a 2D array of (11,11) corresponding to the sum of all possible values of A and B?

For each value of (A+B), the value of C is immediately determined as 1-(A+B), or it does not exist if (A+B)>1. Thus, for each entry in the matrix, you can read off A and B from the column and row labels, and C from 1-(A+B).

Good luck, Craig

```
aa = lindgen(11) # (lonarr(11)+1) ;; A-only matrix
bb = (lonarr(11)+1) # lindgen(11) ;; B-only matrix
cc = 10-(AA+BB)
wh = where(cc GE 0)
a = wh MOD 11 & b = wh/11 & c = cc(wh)
```

Craig B. Markwardt, Ph.D. EMAIL: craigmnet@REMOVEcow.physics.wisc.edu Astrophysics, IDL, Finance, Derivatives | Remove "net" for better response

-----