
Subject: Re: The spectra of fluctuations in variable
Posted by [Craig Markwardt](#) on Thu, 15 Mar 2007 15:20:24 GMT
[View Forum Message](#) <> [Reply to Message](#)

"duxiyu@gmail.com" <duxiyu@gmail.com> writes:

- > I have a time series B(T).
- > The time resolution dt of the data is 4s and the amount N is 1000.
- > The unit of B and T is 'nT' and 's'.
- > I want to study its fluctuations in frequency space using the power
- > spectral density (PSD).
- >
- > BB=FFT(B)
- > freq=FINDGEN(N)/(N*dt)
- >
- > Then I am not clear how to calculate the PSD.
- > Maybe, $PSD=(ABS(BB))^2$, but there is a problem in the unit.
- > The unit of BB is 'nT' that is the same as B, so the unit of PSD which
- > is obtained by the above method is 'nT^2'.
- > But in physical view the unit of PSD should be 'nT^2/Hz'.

If you want "per Hertz" then you need to divide the powers by the spacing between adjacent frequencies.

Also, you will have to worry some about the normalization of the FFT.
Experiment with $FFT(B,+1)$ and $FFT(B,-1)$ until you get what you want.

- > The other question is how to calculate the phase difference between two
- > B1 and B2.
- > I calculate it by the following method, but I am not sure whether it is
- > right.
- >
- > BB1=FFT(B1)
- > BB2=FFT(B2)
- > phase1=ATAN(BB1,/phase)
- > phase2=ATAN(BB2,/phase)
- > difference=phase1-phase2
- >
- > Moreover, the angles seems not smoothed by regular method because of
- > their periodicity.
- > Is there other method to smooth the angle data?

I would normally recommend multiplying the fourier coefficients
to form the "cross spectrum,"

$CR12 = BB1 * CONJ(BB2)$

You can rebin or smooth these complex numbers as you wish, and then
compute ATAN() on the result. Finding the uncertainties of those

values is a much more complicated process.

Good luck,
Craig

--

Craig B. Markwardt, Ph.D. EMAIL: craigmnet@REMOVEcow.physics.wisc.edu
Astrophysics, IDL, Finance, Derivatives | Remove "net" for better response
