Subject: Re: Calculating cumulative probability using cgHistoPlot Posted by David Fanning on Tue, 08 Nov 2011 19:05:13 GMT

View Forum Message <> Reply to Message

Xin Tao writes:

```
> I'm using cghistoplot, and I'm confused by the cumulative probability
> calculated by cghistoplot. I'm wondering whether any one can give me
> some help here.
>
> Suppose we have histdata like this: [2, 0, 0, 3, 5, 4], then if
> we use the way of constoplot to calculate the cumulative probability
> like this:
>
    cumTotal = Total(histData, /CUMULATIVE) ;; gives us
  [2.00000,
               2.00000.
                            2.00000.
                                         5.00000.
                                                     10.0000.
> 14.00001
    probability = Scale_Vector(cumTotal, 0, 1)
                                                  ;; gives us
  probability = [0.00000]
                            0.00000
                                        0.00000
                                                   0.250000
> 0.666667
               1.000001
> This is kind of counter-intuitive to me, because the first value of
> histdata is clearly 2, but the probability is 0 until the 4th value.
> However, I'm not experienced in data anlaysis, and I might have
> misunderstood something about "cumulative probability" here.
>
> It seems to be more natural to me to define the cumulative probability
> in the following way:
>
 probability = total(double(hist)/total(double(hist)), /cumula)
> Am I right?
Probably. I was cribbing some notes from an R class that
I didn't do all that well in. :-(
I'll look into it. :-)
Cheers,
David
David Fanning, Ph.D.
Fanning Software Consulting, Inc.
Coyote's Guide to IDL Programming: http://www.idlcoyote.com/
```

Sepore ma de ni thui. ("Perhaps thou speakest truth.")

Page 2 of 2 ---- Generated from comp.lang.idl-pvwave archive