
Subject: Re: Relative distances between vector elements and search for matches in other vector.

Posted by [Tito](#) on Fri, 27 Apr 2012 14:26:00 GMT

[View Forum Message](#) <> [Reply to Message](#)

Dear Gianguido,

> In general, I am still confused about what you are trying to do. Could you give a simple example of input and output? Something like

> ;inputs:
> r1=[1,3,3,4,5]
> r2=[2,-1,4,3]

I will try to explain :)

I have this vectors:

Wavelength: (lets call it 'spec = Double[40]')

1572.6780	1573.0127	1573.3506	1573.6405	1573.6990
1573.8116	1574.0538	1574.2353	1574.4169	1574.7384
1574.7856	1574.8598	1575.0498	1575.1705	1575.2022
1575.5655	1575.7488	1575.9235	1575.9886	1576.0250
1576.1455	1576.5514	1576.7996	1576.9631	1577.1086
1577.2545	1577.3844	1577.6127	1577.7395	1577.8119
1577.9117	1578.0769	1578.2476	1578.6911	1579.1422
1579.3361	1579.5977	1579.6965	1579.8723	1580.0659

and :

Depths (lets call it 'dep = Double[40]' which I am going to need later on)

0.37406952	0.24975741	0.36841672	0.72394320	0.39689800
0.85735723	0.45238194	0.88476935	0.47967704	0.57341885
0.69364480	0.71783461	0.90553347	0.85711051	0.82898679
0.45230935	0.79680740	0.81328328	0.86826340	0.89866642
0.63804183	0.50215458	0.70966639	0.44995926	0.77566860
0.46117208	0.44478282	0.56285968	0.69161155	0.44447514
0.89239487	0.63570633	0.40267729	0.39813661	0.36184048
0.88953285	0.36669691	0.88203417	0.86033273	0.45937277

So this is All the peaks (actually minimums) of a spectrum. I have Depths on Y axis and Wavelengths on X axis.

Now there is two types of lines that I want to distinguish from 2 catalogs.

Firs one is:

HITRAN = Double[18] wavelengths:

1572.6598	1572.9920	1573.3317	1573.6790	1574.0339
1574.3965	1574.7667	1575.7256	1576.1226	1576.5273

1576.9396	1577.3596	1577.7872	1578.2224	1578.6653
1579.1158	1579.5739	1580.0395		

and the second:

VALD = Double[44] wavelengths:

1573.5713	1573.7811	1573.8207	1574.5017	1574.6219
1574.7876	1574.7883	1575.3189	1575.3291	1575.3502
1575.3588	1575.6021	1575.7971	1576.5622	1576.8632
1576.8819	1576.9953	1577.0055	1577.0098	1577.0150
1577.1689	1577.1878	1577.3403	1577.3732	1577.4928
1577.8381	1578.1040	1578.5977	1578.8859	1578.9233
1579.0978	1579.1627	1579.3312	1579.3826	1580.1757
1580.2550	1580.2879	1580.3051	1580.3843	1580.5266
1580.8015	1581.0597	1581.2374	1581.4767	

Now My idea is to find the distances between the lines in the spectrum and in the catalogs and then identify both populations based on their relative distances.

Thanks to David I did this:

;Finding the relative distance bettwen the HITRAN lines (nm):

```
ratio_hitran = HITRAN[0] - HITRAN
```

;Finding the relative distance bettwen each line on the spectrum:

```
ratio_spec = spec[0] - spec
```

;Finding the relative distance bettwen the lines from the VALD catalog:

```
ratio_vald = VALD[0] - VALD
```

And I get this after substracting:

```
ratio_hitran = 0.0000000  -0.33218057  -0.67191942  -1.0192411  -1.3741670
-1.7367149  -2.1069002  -3.0658462  -3.4628346  -3.8674891
-4.2798071  -4.6997833  -5.1274089  -5.5626720  -6.0055574
-6.4560465  -6.9141175  -7.3797447
```

```
ratio_spec = 0.0000000  -0.33465711  -0.67260343  -0.96251210  -1.0209528
-1.1335506  -1.3758041  -1.5572456  -1.7388401  -2.0604169
-2.1075366  -2.1817349  -2.3717436  -2.4924497  -2.5241943
-2.8874598  -3.0707372  -3.2454473  -3.3105429  -3.3469536
-3.4675066  -3.8733788  -4.1215987  -4.2850644  -4.4306307
-4.5765029  -4.7063881  -4.9347199  -5.0614689  -5.1339078
-5.2336548  -5.3988798  -5.5696123  -6.0131190  -6.4641985
-6.6580319  -6.9196382  -7.0184672  -7.1942482  -7.3878522
```

```
ratio_vald = 0.0000000  -0.20975704  -0.24936781  -0.93035298  -1.0505857
```

-1.2162307	-1.2169309	-1.7475752	-1.7577780	-1.7788837
-1.7874861	-2.0307522	-2.2257052	-2.9908133	-3.2918951
-3.3106002	-3.4239310	-3.4341338	-3.4384350	-3.4436364
-3.5975783	-3.6164834	-3.7689249	-3.8018338	-3.9214663
-4.2667602	-4.5326325	-5.0263668	-5.3145451	-5.3519353
-5.5265028	-5.5913204	-5.7598662	-5.8112802	-6.6043959
-6.6836174	-6.7165264	-6.7337311	-6.8129526	-6.9552913
-7.2301661	-7.4883363	-7.6660846	-7.9053497	

Now in 'ratio_spec' Some of the relative distances belong to ratio_HITRAN and the rest that are left belong to ratio_VALD(some of them).

Example:

ratio_hitran = 0.0000000	-0.33218057	-0.67191942	-1.0192411	-1.3741670
ratio_spec = 0.0000000	-0.33465711	-0.67260343	-0.9625121	-1.3758041

Are exactly the same lines from the spectra and from catalog(they differ slightly in the +/- 0.05 criteria) but I can identify them based on their distance from the first line.

I Want to do it not only respect to the first line, I want to find lines based on their distances one respect to another (example: ratio_spec[i] - ratio_spec[i+1] is the same like ratio_vald[k] - ratio_vald[k+4])

Then also tricky is after I identify the catalog lines in to the spectra, to pass them the depths from the spectra. e.g. where I have match on

spec[0:3]:

1572.6780	1573.0127	1573.3506	1573.6405
-----------	-----------	-----------	-----------

HITRAN[0:3]:

1572.6598	1572.9920	1573.3317	1573.6790
-----------	-----------	-----------	-----------

to give both:

dep[0:3]

0.37406952	0.24975741	0.36841672	0.72394320
------------	------------	------------	------------

and I am able to perform the first step, I am identifying the HITRAN lines and pass them the depts. But this is due to the fact that the first line in 'hitran' and on 'spec' is the same and it is easy. but if this is not the case as it is with the VALD catalog I have no idea how to do it and to pass the depths...

I hope I explain it good.

All the best,
Trifon
