

---

Subject: Re: Coadd images that contain no stars  
Posted by on Wed, 30 May 2012 11:17:57 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Den onsdagen den 30:e maj 2012 kl. 10:58:53 UTC+2 skrev Helder:

>> [http://en.wikipedia.org/wiki/Phase\\_correlation](http://en.wikipedia.org/wiki/Phase_correlation)

>  
> Hi Russel and IDLers,  
> the phase correlation method basically calculates the correlation by using FFT() operations:  
> FFT\_ImgA = FFT(ImgA,-1)  
> FFT\_ImgB = FFT(ImgB,-1)  
> FFT\_ImgA\*CONJ(FFT\_ImgB) / ABS(FFT\_ImgA\*CONJ(FFT\_ImgB))  
> and then locating the maximum (with an appropriate function for higher accuracy).

>  
> On the other hand the Astro-Library Correl\_Images function "computes the 2-D cross-correlation function of two images for a range of (x,y) shifting by pixels of one image relative to the other". This is done using FOR loops for the pixel shifting.

> This is generally speaking not the "IDL-way", but I guess there is no direct way around this. As far as I could see, the Astro I've looked a bit into this and found that the IDL CORREL() function is quite similar but not identical (but maybe I used the wrong keywords...).

>  
> Now the question: how do these two methods compare? FFT procedures will perform well for images with sizes of  $2^n$ , whereas the Astro-Library uses a recursive method to "zoom" in and this might give better accuracy.

>  
> Does anybody have a better idea about how these two methods? When should they be used and when should they not be used?

There are a couple of issues related to the wrap-around caused by the FFT assumption that the image is just one tile in an infinite mosaic of identical tiles. The magnitude of the problems depend on the kind of scene present in your image and the size of the image compared to the expected shifts. If the images are large compared to the expected shifts, the FFT method should be safe and accurate.

Because of the FFT wrap-around, this happens/can happen if your images are small:

- \* The peak corresponding to the actual shift is attenuated because part of the scene is not matched to features in the same "tile" but to features being wrapped in from the other side of the image. This effect is greater the larger the shift really is. It might be negligible if the scene consists mainly of some small but bright features entirely within the field of view (FOV) and more serious if the scene is extended with features that are about as strong over the entire FOV.

- \* The correlation peak can also be attenuated if the matching feature is small in extent compared to the pixels and fall differently on the pixel grid in one image compared to the other.

- \* There may be false peaks caused by features on one side of the FOV being similar to features wrapping in from the other side. These false peaks should normally be weaker than the peak corresponding to the true match but if the true peak is attenuated by the effect explained above, the false peak may be higher. (In most situations the risk that this happens should be small.)

- \* The wrapping in of features from the other side of the image also affects the `_shape_` of the

correlation peak because the side of the peak corresponding to too large shifts is more attenuated than the side corresponding to too small shifts. If you are interested in interpolation of the peak in order to reach subpixel accuracy, this effect moves the peak toward smaller shifts and therefore causes a small but systematic underestimation of the shifts.

Calculating your correlation function in the image domain is safe from these effects also for small images (although you have to sacrifice some of the FOV so you evaluate the correlation also for the largest considered shifts with actual scene within the evaluation window for both images). In addition you can consider other metrics than correlation for the best match.

I looked into matters of this sort a couple of years ago, there is a paper in A&A that you might find useful: <http://adsabs.harvard.edu/abs/2010A%26A...524A..90L>

I had a particular application in mind of course, but some of the methods and reasoning should be relevant to other situations. In addition to some methods used for calculating the correlation peak, I also discuss different methods for interpolating the peak in order to get shifts with subpixel accuracy.

---