

---

Subject: Re: Gradient of two dimensional field  
Posted by [Mirko Vukovic](#) on Wed, 19 Feb 1997 08:00:00 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Andy Loughe wrote:

>  
> Wilpert\_Martin wrote:  
>>  
>> Hi everybody,  
>>  
>> we want to determine the electrical field from a given potential,  
>> i.e. we have to calculate the gradient of a two dimensional array.  
>>  
>> Has anybody a idl-pvwave procedure to do this task?  
>>  
>> Thank you in advance  
>>       Martin Wilpert  
>  
> I would think that the shift function (used twice)  
> could be used to do this.  
>  
> --  
> Andrew F. Loughe                   |  
> afl@cdc.noaa.gov  
> University of Colorado, CIRES Box 449 |  
> http://cdc.noaa.gov/~afl  
> Boulder, CO 80309-0449           | phn:(303)492-0707  
> fax:(303)497-7013  
> -----  
> "I do not feel obliged to believe that the same God who has endowed us  
> with  
> sense, reason, and intellect has intended us to forego their use."  
> -Galileo  
or, convolve with a kernel that will give you the gradient.  
(we are talking two kernels, one for each direction)  
--

Mirko Vukovic, Ph.D   3075 Hansen Way M/S K-109  
Varian Associates Palo Alto, CA, 94304  
415/424-4969   mirko.vukovic@varian.grc.com

---