
Subject: Re: Generating N random numbers that add to a TOTAL

Posted by [Russell Ryan](#) on Thu, 07 Aug 2014 14:08:03 GMT

[View Forum Message](#) <> [Reply to Message](#)

On Wednesday, August 6, 2014 11:52:47 PM UTC-4, Gianguido Cianci wrote:

> Hi all,

>

>

>

> I am wondering if anybody has suggestions on how to improve the function below. It seems ok for floating precision numbers.

>

>

>

> For integers it's a different story:

>

> It works great if $N \ll \text{TOTAL}$. When N approaches TOTAL I get a few numbers and then a bunch of zeros... Also, setting /DIFFERENT makes it run for ever if N is large. Also, the sum of res adds up $\text{TOTAL} \pm 1$, not always to TOTAL exactly...

>

>

>

> Suggestions?

>

>

>

> Thanks,

>

> Gianguido

>

>

>

>

>

>

>

> FUNCTION nrndaddto, n, total, integers = integers, different = different

>

>

>

> compile_opt idl2

>

>

>

> res = dblarr(n)

>

> res[0] = randomu(seed, 1, /double)*(total)

>

```

>
>
> FOR i = 1, n-2 DO BEGIN
>
>   res[i] = randomu(seed, 1, /double)*(total-total(res[0:i-1], /double))
>
> ENDFOR
>
> res[n-1] = total-total(res[0:n-2], /double)
>
>
>
> IF ~keyword_set(integers) THEN integers = 0
>
>
>
> IF keyword_set(integers) THEN res = round(res)
>
> IF keyword_set(different) THEN BEGIN
>
>   IF n_elements(res) NE n_elements(unique(res, /sort)) THEN res = $
>
>   nrndaddto(n, total, integers = integers, different = 1)
>
> ENDIF
>
>
>
>
> RETURN, res
>
> END

```

I should've said. Since the running total is converging to the requested total, the numbers are getting smaller with time. That is almost certainly related to why your sequence doesn't seem uniform.

R
