
Subject: Re: Complex -> Real FFT

Posted by [Jeff Wasserman](#) on Mon, 26 Jul 1999 07:00:00 GMT

[View Forum Message](#) <> [Reply to Message](#)

David Fanning wrote:

>
> Jeremy Kepner (jvkepner@muse.Princeton.EDU) writes:
>
>> Does anyone know of a Complex to Real FFT that has
>> been implemented in IDL. (Likewise a corresponding
>> Real to Complex FFT would also be nice).
>
> I'm probably missing something deep here, but
> doesn't this do what you want?
>
> result = FLoat(complex_fft_expression)
> result = Imaginary(complex_fft_expression)

If you don't care about phase information with your FFT data, and instead want just an absolute power measurement (like a spectrum analyzer), you could use

```
power_spectrum = Abs(complex_fft_data)
```

[the abs function returns the value of $\sqrt{\text{real}^2 + \text{imaginary}^2}$ if it's argument is complex]

Conversely, you could also get the phase info into an array

```
phase_array = Tan(complex_fft_data)
```

[The tan function also nicely works with complex data]

From these two arrays you could then go back to the complex array. However, as far as I know, there is no way to obtain the two independent variables of the original complex array after it's been converted to reals.

Jeffrey Wasserman - Member of Technical Staff
Group 83 Electro-Optical Materials and Devices
MIT Lincoln Laboratory Lexington, MA

Warning - I am using a government mail server, any unauthorized spam mail will be prosecuted! Change the x's to dots and the y to 'at' sign to respond: wassermany@llxmit.edu
