
Subject: Re: PLOT question

Posted by [davidf](#) on Wed, 20 Dec 2000 18:16:41 GMT

[View Forum Message](#) <> [Reply to Message](#)

Stuart Colley (src@star.ucl.ac.uk) writes:

- > I have some data that is periodic, $y=f(t)$ and the data is plotted as
- > plot, t, y The problem is I'd like t to plot phase angle rather than
- > time, the problem being the phase angle is between 0 and 1 and repeats.
- > Attempting to plot the data you just get a plot between 0 and 1, what I'd
- > like to be able to plot is each cycle of the data with it's phase angle.
- >
- > Any ideas on how to get the plot axes right?

If you are just looking for ideas, why not use the plot you have but add the phase angle as a symbol on the plot? You could have a filled circle for a phase angle of 1, a half-filled circle for a phase angle of 0.5, etc.

Shouldn't be too hard to come up with something like that. :-)

Cheers,

David

--

David Fanning, Ph.D.

Fanning Software Consulting

Phone: 970-221-0438 E-Mail: davidf@dfanning.com

Coyote's Guide to IDL Programming: <http://www.dfanning.com/>

Toll-Free IDL Book Orders: 1-888-461-0155

Subject: Re: PLOT question

Posted by [Dave Greenwood](#) on Wed, 20 Dec 2000 18:32:54 GMT

[View Forum Message](#) <> [Reply to Message](#)

Stuart Colley <src@star.ucl.ac.uk> wrote:

- >
- > I have some data that is periodic, $y=f(t)$ and the data is plotted as
- > plot, t, y The problem is I'd like t to plot phase angle rather than
- > time, the problem being the phase angle is between 0 and 1 and repeats.
- > Attempting to plot the data you just get a plot between 0 and 1, what I'd
- > like to be able to plot is each cycle of the data with it's phase angle.
- >
- > Any ideas on how to get the plot axes right?

If I understand your question correctly and assuming that the phase angle is a function of t, something like the following might work:

```
FUNCTION PHASE_ANGLE, axis, index, t
  RETURN, STRING(phase angle at time t)
END
```

..

```
PLOT, T, Y, XTICKFORMAT='PHASE_ANGLE'
```

Dave

Dave Greenwood Email: Greenwoodde@ORNL.GOV
Oak Ridge National Lab %STD-W-DISCLAIMER, I only speak for myself

Subject: Re: PLOT question

Posted by [Paul van Delst](#) on Wed, 20 Dec 2000 18:51:24 GMT

[View Forum Message](#) <> [Reply to Message](#)

Stuart Colley wrote:

>

> I have some data that is periodic, $y=f(t)$ and the data is plotted as
> plot, t, y The problem is I'd like t to plot phase angle rather than
> time, the problem being the phase angle is between 0 and 1 and repeats.
> Attempting to plot the data you just get a plot between 0 and 1, what I'd
> like to be able to plot is each cycle of the data with it's phase angle.

>

> Any ideas on how to get the plot axes right?

If I understand you correctly, you want to remove the discontinuity in your phase angle - like in a TAN plot that goes from $\pm \pi$ or $\pm \pi/2$?

If so, one way is to calculate the phase angle derivatives. If there is a change in sign near a known boundary (0 or 1), you can add or subtract (based on the sign of the derivative) the required amount to make the phase angle curve continuous.

I did this many years ago in fortan to view the phase of radiometric signals from an interferometer. They bounced between $\pm \pi$ and doing the above made the plot relatively (insert some hand-waving here) continuous. If you have a noisy signal, then it can get a bit more difficult as the noise can confuse matters when the signal is close to a boundary, i.e. you get a lot of flipping back and forth, but then the result is a continuous curve with dropouts which, in some respects, may also be a useful plot.

If this is not what you mean, then delete this message.

paulv.

--

Paul van Delst Ph: (301) 763-8000 x7274
CIMSS @ NOAA/NCEP Fax: (301) 763-8545
Rm.207, 5200 Auth Rd. Email: pvandelst@ncep.noaa.gov
Camp Springs MD 20746

Subject: Re: PLOT question

Posted by [Craig Markwardt](#) on Thu, 21 Dec 2000 00:01:42 GMT

[View Forum Message](#) <> [Reply to Message](#)

Stuart Colley <src@star.ucl.ac.uk> writes:

- > I have some data that is periodic, $y=f(t)$ and the data is plotted as
- > plot, t, y The problem is I'd like t to plot phase angle rather than
- > time, the problem being the phase angle is between 0 and 1 and repeats.
- > Attempting to plot the data you just get a plot between 0 and 1, what I'd
- > like to be able to plot is each cycle of the data with it's phase angle.
- >
- > Any ideas on how to get the plot axes right?

It's not clear to me that everybody else is going waaaayyy overboard here. Stuart hasn't responded so that may be a good or a bad sign.

Are you just asking how to convert linear time into phase? Perhaps this is best accomplished with the MOD operator. If your signal has period P, and the starting epoch is T0, then this may do the trick.

```
phase = (t-t0)/p MOD 1
phase = (phase+1) MOD 1
plot, phase, y
```

The "MOD 1" takes the remainder when divided by 1, or simply the fractional value. The "extra" MOD step there is because MOD doesn't handle negative numbers very well. It maps the segment from -1 to 0 back onto 0 to 1. Another way to do it in one step is

```
phase = (t-t0)/p
phase = phase - floor(phase)
```

Craig

--

Craig B. Markwardt, Ph.D. EMAIL: craigmnet@cow.physics.wisc.edu
Astrophysics, IDL, Finance, Derivatives | Remove "net" for better response

Subject: Re: PLOT question
Posted by [Craig Markwardt](#) on Thu, 21 Dec 2000 00:14:52 GMT
[View Forum Message](#) <> [Reply to Message](#)

Craig Markwardt <craigmnet@cow.physics.wisc.edu> writes:

> Stuart Colley <src@star.ucl.ac.uk> writes:
>
>> I have some data that is periodic, $y=f(t)$ and the data is plotted as
>> plot, t, y The problem is I'd like t to plot phase angle rather than
>> time, the problem being the phase angle is between 0 and 1 and repeats.
>> Attempting to plot the data you just get a plot between 0 and 1, what I'd
>> like to be able to plot is each cycle of the data with it's phase angle.
>>
>> Any ideas on how to get the plot axes right?
>

Oops now that I look at the question again, I think maybe Stuart was just looking for $(T-T_0)/P$, without the MOD? Or, a way to sandwich together many distant cycles? If the latter, that's a bit more difficult. A small program would be needed to join these intervals together.

Craig

--

Craig B. Markwardt, Ph.D. EMAIL: craigmnet@cow.physics.wisc.edu
Astrophysics, IDL, Finance, Derivatives | Remove "net" for better response

Subject: Re: PLOT question
Posted by [davidf](#) on Thu, 21 Dec 2000 00:40:19 GMT
[View Forum Message](#) <> [Reply to Message](#)

Craig Markwardt (craigmnet@cow.physics.wisc.edu) writes:

> Oops now that I look at the question again, I think maybe Stuart was
> just looking for $(T-T_0)/P$, without the MOD? Or, a way to sandwich

- > together many distant cycles? If the latter, that's a bit more
- > difficult. A small program would be needed to join these intervals
- > together.

Oh, now, writing a *program* is WAY over the top!

Cheers,

David

P.S. Let's just say this is the reason I only suggested an *idea*. :-)

--

David Fanning, Ph.D.

Fanning Software Consulting

Phone: 970-221-0438 E-Mail: davidf@dfanning.com

Coyote's Guide to IDL Programming: <http://www.dfanning.com/>


Toll-Free IDL Book Orders: 1-888-461-0155

Subject: Re: PLOT question


Posted by [Stuart Colley](#) on Thu, 21 Dec 2000 11:46:32 GMT

[View Forum Message](#) <> [Reply to Message](#)

Maybe a small example will be of help - hopefully it will be a bit clearer than my first message. Imagine a periodic signal (sin wave for example) and we are plotting more than one cycle, Normally the signal would be plotted as a function of time, e.g:



What I want to do is swap time for phase angle, so the phase is somewhere between 0 - 2pi, but the problem is that the phase angle repeats, so what I was looking for on the axis was something like:



|
end of first cycle, start of second

What I'll probably do is use `xtickv` and `xtickname` to set the ticks, since as far as I can tell, you MUST 'plot, t, y' to get the plot right, 'plot, phase, y' results in the data being merged together and only one cycle being plotted.

The reason I sent the first post - I was wondering if there was an easier way than `xtickv/xtickname` since it can't automatically choose the tick values like PLOT can, and isn't so flexible when I'm trying to write a general purpose routine. Time for some more code I think....

cheers,
S

Subject: Re: PLOT question

Posted by [Jaco van Gorkom](#) on Thu, 21 Dec 2000 14:05:40 GMT

[View Forum Message](#) <> [Reply to Message](#)

Stuart Colley wrote:

> What I want to do is swap time for phase angle, so the phase is
> somewhere between 0 - 2pi, but the problem is that the phase angle
> repeats, so what I was looking for on the axis was something
> like:

>
> y value |
> |
> | x x
> | x x x x x
> _____ x _____ x _____
> time | 0 pi/2 pi 3pi/2 0 pi/2 pi 3pi/2 0 etc...
>

>
> |
> end of first cycle, start of second
>

> What I'll probably do is use `xtickv` and `xtickname` to set the ticks, since
> as far as I can tell, you MUST 'plot, t, y' to get the plot right,
> 'plot, phase, y' results in the data being merged together and only one
> cycle being plotted.

>
> The reason I sent the first post - I was wondering if there was an easier
> way than `xtickv/xtickname` since it can't automatically choose the tick
> values like PLOT can, and isn't so flexible when I'm trying to write a
> general purpose routine.

The XTICKFORMAT mechanism as proposed by Dave should give you the periodic phase angle tick values, if you plug in the logic by Craig:

```
FUNCTION PHASE_ANGLE, axis, index, t
  phase = (t-t0)/period
  phase = phase - floor(phase)
  ; this phase angle goes from 0 to 1, you want 0 - 2pi:
  phase = 2*!PI*phase
  RETURN, STRING(phase)
END

..
PLOT, t, y, XTICKFORMAT='PHASE_ANGLE'
```

Now PLOT will automatically choose "nice" tick values based on the original time values, e.g. 1 s, 2 s, ... You would like "nice" values in phase angle (pi, 2pi, ..), so I would plot as a function of t/period (actually the total phase angle), just to get the tick marks at nice locations:

```
FUNCTION PHASE_ANGLE, axis, index, totalphase
  phase = totalphase - floor(totalphase)
  ; this phase angle goes from 0 to 1, you want 0 - 2pi:
  phase = 2 * !pi * phase
  RETURN, (FORMAT_AXIS_VALUES(phase))[0]
END

..
t0 = 0 ; starting time of first (or any) period
PLOT, (t-t0)/period, y, XTICKFORMAT='PHASE_ANGLE'
```

One important thing here: you need to make sure that IDL uses at least two tick marks per period, or else these nice periodic tick values will look just like a ordinary series of zeroes. (for this, set the XTICKS keyword if necessary)

I used FORMAT_AXIS_VALUES to get "nice" formatting of the strings. If you *really* want "3pi/2" at your tick mark instead of 4.71, then that should not be too difficult to code into the PHASE_ANGLE function. Better left as an exercise to the reader...

Jaco

Jaco van Gorkom gorkom@rijnh.nl
FOM-Instituut voor Plasmafysica Rijnhuizen

Subject: Re: PLOT question

[View Forum Message](#) <> [Reply to Message](#)

V V V V V V

Cheers,
Martin

—

```

[[ Dr. Martin Schultz   Max-Planck-Institut fuer Meteorologie [[
[[           Bundesstr. 55, 20146 Hamburg                      [[
[[           phone: +49 40 41173-308                            [[
[[           fax: +49 40 41173-298                               [[
[[ martin.schultz@dkrz.de                                       [[

```