

---

Subject: warping continents to magnetic coordinates  
Posted by [Ken Mankoff](#) on Tue, 22 May 2001 22:55:19 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hi everyone,

I am interested in displaying the continents on a non geographic coordinate grid.

Specifically, I have the corrected geomagnetic CGM grid off the NSSDC website, and wish to display the continents with some data on this grid. Those of you with the latest "Reviews of Geophysics" (Volume 39, Number 2, May 2001) can see on the cover what I'm trying to do.

I have found the maps in the IDL map directory. I have read (briefly) the map\_continents procedure. I discovered the !map system variable. I am just not sure what my next step should be...

Thanks for any help,

-k.

--

Ken Mankoff  
LASP://303.492.3264  
<http://lasp.colorado.edu/~mankoff/>

---

---

Subject: Re: warping continents to magnetic coordinates  
Posted by [Brian Jackel](#) on Wed, 23 May 2001 17:09:14 GMT  
[View Forum Message](#) <> [Reply to Message](#)

---

Hi Ken

I've got some code that you may find useful. It works in "geomagnetic dipole" coordinates as defined in Russell or Hapgood. Fairly similar to CGM in most places.

The basic idea is to pull data directly from the IDL map files, then apply a coordinate transformation. The rotation matrix is obtained from some \*VERY\* rough code of mine called "geospace\_convert". It seems to work well for this purpose, but please use with caution.

Let me know if you have any trouble with this. It \*should\* just compile and run. Again, this is not

high quality code, but hopefully you'll find it useful.

Brian

```
;1999/08/03    Brian Jackel
;
; This IDL script draws a continental map in dipole geomagnetic
coordinates.
; It takes a few key lines from "map_continents" and does coordinate
transformation
; using "geospace_convert".
;

name= FILEPATH('clow.ndx',SUBDIR=['resource','maps','low'])
OPENR,lun,fname,/XDR,/GET_LUN

nsegments=0L
READU,lun,nsegments

result= REPLICATE({fptr:0L, npts:0L, latmax:0.0, latmin:0.0,
lonmax:0.0, lonmin:0.0},nsegments)
READU,lun,result
FREE_LUN,lun

map_set,90,-50,79,/GNOMIC,LIMIT=[55,-120,85,190]
re= 6371.2

;
;

;rotation_matrix= FLTARR(3,3)
;rotation_matrix[0,0]= 1.0
;rotation_matrix[1,1]= 1.0
;rotation_matrix[2,2]= 1.0

CDF_EPOCH,Epoch,1990,1,1,23,0,0,/COMPUTE_EPOCH
dummy=
GEOSPACE_CONVERT([1,1,1],'geo','mag',ROTATION_MATRIX=rotation_matrix,
CDF_EPOCH_DATE=epoch)

fname= FILEPATH('clow.dat',SUBDIR=['resource','maps','low'])
OPENR,lun,fname,/XDR,/GET_LUN

FOR indx=0,nsegments-1 DO BEGIN
  IF (result[indx].npts GE 2) THEN BEGIN
    POINT_LUN,lun,result[indx].fptr
```

```

xy= FLTARR(2,result[indx].npts,/NOZERO)
READU,lun,xy

R= 1
lat_rad= REFORM(xy[0,*]) *!dtr
lon_rad= REFORM(xy[1,*]) *!dtr
temp= FLTARR(result[indx].npts,3)
temp(0,0)= R * cos(lat_rad) * cos(lon_rad)
temp(0,1)= R * cos(lat_rad) * sin(lon_rad)
temp(0,2)= R * sin(lat_rad)
temp= rotation_matrix###temp

l= SQRT( temp[* ,0]^2 + temp[* ,1]^2 )

xy[0,*]= ATAN(temp[* ,2],l) * !radeg
xy[1,*]= ATAN(temp[* ,1],temp[* ,0]) * !radeg

PLOTS,xy[1,*],xy[0,*],NOCLIP=0,/DATA,;COLOR=128
ENDIF
ENDFOR

FREE_LUN,lun

;!p.font=1
;DEVICE, SET_FONT='Helvetica Bold Italic', /TT_FONT

MAP_GRID,lats=[50,55,60,65,70,75,80,85],lons=[-35]

;Contwoyto Lake  65.75 248.75  72.64 298.10
;Fort Simpson   61.76 238.77  67.27 290.66

;Hopen Island   76.51 25.01  71.92 131.66
;Soroya         70.54 22.22  67.30 120.50

;USERSYMBOL,'star',/FILL,SIZE=2
PLOTS,298.10,72.64,PSYM=5 ;cont
XYOUTS,298.10,72.64,'Contwoyto Lake',CHARSIZE=2
PLOTS,131.66,71.92,PSYM=5 ;hop
XYOUTS,131.66,71.92,'Hopen Island',CHARSIZE=2

USERSYMBOL,'triangle',/FILL,SIZE=2
PLOTS,290.66,67.27,PSYM=5 ;sim
XYOUTS,290.66,67.27,'Fort Simpson',CHARSIZE=2
PLOTS,120.50,67.30,PSYM=5 ;sor
XYOUTS,120.50,67.30,'Soroya',CHARSIZE=2

```

```
;MAP_GRID,lats=[0,90],lons=[298.1,290.66,131.66,120.5],GLINE STYLE=0
```

```
;!p.font=0
```

```
END
```

```
;+
```

```
; NAME:      GEOSPACE_CONVERT
```

```
; PURPOSE:   Conversion between various geocentric Cartesian  
;            space physics coordinate systems.
```

```
; CATEGORY:  Coordinate systems
```

```
; CALLING SEQUENCE:  a= GEOSPACE_CONVERT( Position, From, Into )
```

```
; INPUTS:
```

```
;   Position    a 3 element vector or 3xN element array  
;               containing the position(s) to be converted.  
;               Usually in Cartesian coordinates, use  
;               /FROM_SPHERICAL if in spherical coordinates.
```

```
;   From    the names of the initial and final coordinate  
;   Into    systems. Must be a 2-3 letter string from the set  
;           of 6 allowable coordinate systems:
```

```
;       GEI   Geocentric equatorial inertial  
;       GEO   Geographic (rotating earth)  
;       GSE   Geocentric solar ecliptic  
;       GSM   Geocentric solar magnetospheric  
;       SM    Solar magnetic  
;       MAG   Geomagnetic
```

```
; KEYWORD PARAMETERS: Many of the coordinate systems require time  
;                      information, which can be provided via the  
;                      following keywords. Note that JD starts at  
;                      noon, while UT starts at midnight. Not my idea.
```

```
; JULIAN_DAY    Result = JULDAY(Month, Day, Year)
```

```
; UNIVERSAL_TIME
```

```
; CDF_EPOCH
```

```
; OUTPUTS:      a 3 element vector or 3 x n element array the  
;               same size as input Position. Usually Cartesian,  
;               unless the /INTO_SPHERICAL keyword is set, in
```

```

;           which case it will be (Radius,Latitude,Longitude)
;
;
; RESTRICTIONS:
;
; PROCEDURE:  Construct the appropriate rotation matrices, combine
them
;             and apply to the initial coordinates.  For a detailed
;             description, see:
;
;             Space Physics Coordinate Transformations: A User Guide
;             M. A. Hapgood, Planetary and Space Science,
;             Volume 40, Number 5, pages 711-717, 1992
;
;
; MODIFICATION HISTORY:
; September 8 1993    Written by Brian Jackel, University of Western
Ontario
;
;
;-
;-----

```

```

function GEOSPACE_ROTATION,angle,axis
;
; This is a short utility routine for creating rotation matrices.
;
; zeta is rotation angle in degrees
; axis is rotation axis, a string of 'X','Y' or 'Z'
;
; The basic idea is to
;   start with an empty 3x3 matrix
;   put a 1 in the diagonal element corresponding to the rotation axis
;   put cos(zeta) in the other diagonal elements
;   determine the two off-diagonal terms in the same columns and rows as
the cos(zeta) values
;   put -sin(zeta) in the term above the diagonal
;   put sin(zeta) in the term below
;
;
; element #'s      example for a 'Z' rotation
; 0 1 2          cos(zeta) -sin(zeta)  0
; 3 4 5          sin(zeta)  cos(zeta)  0
; 6 7 8          0          0          1
;
;
sin_zeta= sin(angle!*dtor)
cos_zeta= cos(angle!*dtor)

fill= [1.0d0,cos_zeta,cos_zeta,-sin_zeta,sin_zeta]

```

```

rmatrix= fltarr(3,3)
CASE STRUPCASE(axis) OF
'X': rmatrix([0,4,8,7,5])= fill
'Y': rmatrix([4,0,8,6,2])= fill
'Z': rmatrix([8,0,4,3,1])= fill
ELSE: MESSAGE,'AXIS parameter must be string of X, Y, or Z'
ENDCASE

```

```

RETURN,rmatrix
END

```

```

;-----
function GEOSPACE_CONVERT,Position,From,Into,HELP=help, $
    ROTATION_MATRIX=rotation_matrix, $
    CDF_EPOCH_DATE=cdf_epoch_date, $
    YMDHMS_DATE=ymdhms_date, $
;    JULIAN_DATE=julian_date,
UNIVERSAL_TIME=universal_time, $
    FROM_SPHERICAL=from_spherical,
INTO_SPHERICAL=into_spherical

```

```

;ON_ERROR,2

```

```

IF KEYWORD_SET(HELP) THEN BEGIN
DOC_LIBRARY,'Geospace_Convert'
return, -1.0
END

```

```

IF (N_PARAMS() EQ 0) THEN MESSAGE,"Error- input value Position not
defined"

```

```

siz= SIZE(position)
; IF (siz(0) EQ 0) THEN MESSAGE,"Error- input value Position not
defined"
; IF (siz(0) GT 2) THEN MESSAGE,"Error- Position must be a 3 element
vector, or a 3xn element array"
; IF (siz(1) NE 3) THEN MESSAGE,"Error- Position must be a 3 element
vector, or a 3xn element array"

```

```

_from= STRMID( STRCOMPRESS(STRUPCASE(from),/REMOVE_ALL) ,0,3 )
_into= STRMID( STRCOMPRESS(STRUPCASE(into),/REMOVE_ALL) ,0,3 )
IF (_from EQ _into) THEN RETURN,position ;identity transformation

```

```

;
;
;
;
;

```

```
seconds_in_day= 24.0d0 * 60.0d0 * 60.0d0
seconds_in_century= 36525.0d0 * seconds_in_day
```

```
CDF_EPOCH,epoch2000,2000,01,01,12,/COMPUTE
CDF_EPOCH,epochunix,1970,01,01,/COMPUTE
CDF_EPOCH,epochmjd,1858,11,17,/COMPUTE
```

```
IF KEYWORD_SET(CDF_EPOCH_DATE) THEN BEGIN
```

```
CDF_EPOCH,cdf_epoch_date,year,month,day,hour,minute,second,/ BREAKDOWN
  IF (year LT 1970) OR (year GT 2100) THEN MESSAGE,'Note- year
'+STRING(year)+' is outside expected range',/INFORMATIONAL
```

```
  epoch= cdf_epoch_date
ENDIF ELSE IF KEYWORD_SET(YMDHMS_DATE) THEN BEGIN
  ymdhms= ymdhms_date
  CASE N_ELEMENTS(ymdhms) OF
    1:CDF_EPOCH,epoch,ymdhms[0],/COMPUTE
    2:CDF_EPOCH,epoch,ymdhms[0],ymdhms[1],/COMPUTE
    3:CDF_EPOCH,epoch,ymdhms[0],ymdhms[1],ymdhms[2],/COMPUTE
    4:CDF_EPOCH,epoch,ymdhms[0],ymdhms[1],ymdhms[2],ymdhms[3],/C OMPUTE
```

```
5:CDF_EPOCH,epoch,ymdhms[0],ymdhms[1],ymdhms[2],ymdhms[3],ym dhms[4],/COMPUTE
```

```
6:CDF_EPOCH,epoch,ymdhms[0],ymdhms[1],ymdhms[2],ymdhms[3],ym
dhms[4],ymdhms[5],/COMPUTE
  ELSE:MESSAGE,'Error- keyword parameter YMDHMS must have between 1
(just year) and 6 (year, month, day, hour, minute, second) elements'
```

```
  ENDCASE
ENDIF ELSE BEGIN
  MESSAGE,'Warning- no time provided, using current system
time',/INFORMATIONAL
  st= SYSTIME(1)*1.0d3      ;convert from second to milliseconds
  epoch= epochunix + st
ENDELSE
```

```
; Determine the modified julian date (since 00:00 UT 17 November 1858)
; and time of day in hours. Then calculate "T_0", the time in julian
; centuries from 12:00 UT on 1 January 2000 (epoch 2000) to the previous
; midnight.
```

```
;
mjddate= (epoch - epochmjd)/1d3      ;modified julian date
in seconds
utime= (mjddate MOD seconds_in_day) / 3600.0d0 ;time of day in hours
mjddate= LONG(mjddate / seconds_in_day)      ;modified julian date
in days, truncated to integer
```

```
T_0= (mjdate - 51544.5d0) / 36525.0d0
```

```
; If the keyword is set, convert from spherical to
; Cartesian coordinates. Otherwise just make a
; working copy of the position vector.
;
new_position= position
IF KEYWORD_SET(FROM_SPHERICAL) THEN BEGIN
  R= position(0,*)
  lat_rad= position(1,*) *!dtr
  lon_rad= position(2,*) *!dtr
  new_position(0,0)= R * cos(lat_rad) * cos(lon_rad)
  new_position(1,0)= R * cos(lat_rad) * sin(lon_rad)
  new_position(2,0)= R * sin(lat_rad)
ENDIF
```

```
; namelist= ['SM','GEO','GEI','GSE','GSM','SM']
; tlist= [-5,-1,2,3,4]
; w= WHERE
```

```
;=====
=====
;Determine what transformation ("T") matrices are required. This is
just a 6x6
;lookup table, from which we get a sequence of transformations. The
number indicates
;the transformation matrix (using Hapgood's convention), with 0 just a
place keeper.
;
CASE _from OF
  'GEI': matrices= [ [0,0,0], [1,0,0], [2,0,0],
[3,2,0], [4,3,2], [5,1,0] ]
  'GEO': matrices= [ [-1,0,0,0], [0,0,0,0], [2,-1,0,0],
[3,2,-1,0], [4,3,2,-1], [5,0,0,0] ]
  'GSE': matrices= [ [-2,0,0], [1,-2,0], [0,0,0],
[3,0,0], [4,3,0], [5,1,-2] ]
  'GSM': matrices= [ [-2,-3,0,0], [1,-2,-3,0], [-3,0,0,0],
[0,0,0,0], [4,0,0,0], [5,1,-2,-3] ]
  'SM': matrices= [ [-2,-3,-4,0,0], [1,-2,-3,-4,0], [-3,-4,0,0,0],
[-4,0,0,0,0], [0,0,0,0,0], [5,1,-2,-3,-4] ]
  'MAG': matrices= [ [-1,-5,0,0,0], [-5,0,0,0,0], [2,-1,-5,0,0],
[3,2,-1,-5,0], [4,3,2,-1,-5], [0,0,0,0,0] ]
ELSE:MESSAGE,'Parameter From must be one of GEI, GEO, GSE, GSM, SM,
MAG'
```

ENDCASE

CASE \_into OF

'GEI': matrices= REFORM( matrices(\*,0) )

'GEO': matrices= REFORM( matrices(\*,1) )

'GSE': matrices= REFORM( matrices(\*,2) )

'GSM': matrices= REFORM( matrices(\*,3) )

'SM': matrices= REFORM( matrices(\*,4) )

'MAG': matrices= REFORM( matrices(\*,5) )

ELSE:MESSAGE,'Parameter Into must be one of GEI, GEO, GSE, GSM, SM,  
MAG'

ENDCASE

matrices= matrices( WHERE(matrices NE 0) ) ;throw away any  
zeros

;-----

;=====

=====

;This is where any required rotation matrices are actually constructed.

;Make a stack of 6 3x3 matrices, but only calculate the necessary ones.

;Note that some matrices may require partial or full evaluation of

;other matrices, so group some calculations together depending on what

;is required. The result is a little cryptic, but reasonably efficient.

;

;Make a stack for 6 3x3 rotation matrices, only the top five of which  
will be used.

;Leave the 0th one empty to keep consistent with notation in Hapgood

tstack= FLTARR(3,3,6)

w1= TOTAL( ABS(matrices) EQ 1 )

w2= TOTAL( ABS(matrices) EQ 2 )

w3= TOTAL( ABS(matrices) EQ 3 )

w4= TOTAL( ABS(matrices) EQ 4 )

w5= TOTAL( ABS(matrices) EQ 5 )

IF (w1 OR w3 OR w4) THEN BEGIN

theta= 100.461d0 + 36000.770d0\*T\_0 + 15.04107d0\*utime

;Greenwich mean sidereal time

tstack(0,0,1) = GEOSPACE\_ROTATION(theta,'Z') ;"T1" in Hapgood,  
equation 2

ENDIF

IF (w2 OR w3 OR w4) THEN BEGIN

ecliptic\_obliquity= 23.439d0 - 0.013d0\*T\_0

mean\_anomaly= 357.528d0 + 35999.050d0\*T\_0 + 0.04107d0\*utime

```

    mean_longitude= 280.460d0 + 36000.772d0*T_0 + 0.04107d0*utime
    ecliptic_longitude= mean_longitude + (1.915d0 -
0.0048d0*T_0)*sin(mean_anomaly!*dtor) + 0.020*sin(2*mean_anomaly!*dtor)
    Ez= GEOSPACE_ROTATION(ecliptic_longitude,'Z')
    Ex= GEOSPACE_ROTATION(ecliptic_obliquity,'X')
    tstack(0,0,2) = Ez##Ex
ENDIF

IF (w3 or w4 or w5) THEN BEGIN

    dyear= (mjdate-46066.0d0)/365.25    ;should be centered on 1985
    phi= 78.7646 + 0.0351936*dyear -0.000128034*dyear^2
    lambda= 289.097 - 0.0363523*dyear -0.00142802*dyear^2

    IF (w5) THEN BEGIN
        Ez= GEOSPACE_ROTATION(lambda,'Z')
        Ey= GEOSPACE_ROTATION(phi-90.0,'Y')
        tstack(0,0,5) = Ey##Ez
    ENDIF

ENDIF

IF (w3 or w4) THEN BEGIN
    phi_rad= phi!*dtor
    cos_phi_rad= cos(phi_rad)
    lambda_rad= lambda!*dtor
    Qg= [ cos_phi_rad*cos(lambda_rad), cos_phi_rad*sin(lambda_rad),
sin(phi_rad) ]
    Qe= REFORM(tstack[*,* ,2]) ## ( TRANSPOSE(REFORM(tstack[*,* ,1]))##Qg
) ;equation (7)

    IF (w3) THEN BEGIN
        psi= ATAN( Qe(1),Qe(2) )!*radeg
        tstack(0,0,3) = GEOSPACE_ROTATION(-psi,'X')
    ENDIF

    IF (w4) THEN BEGIN
        mu= ATAN( Qe(0), SQRT( Qe(1)^2+Qe(2)^2 ) )!*radeg
        tstack(0,0,4)= GEOSPACE_ROTATION(-mu,'Y')
    ENDIF

ENDIF

;
;
;Combine the "T" matrices, do inversions (transpositions) on
;those matrices that need it
;
nstack= N_ELEMENTS(matrices)

```

```

T= fltarr(3,3)
T([0,4,8])= [1.0,1.0,1.0]      ;set diagonal elements equal to 1
(identity matrix)
FOR i=nstack-1,0,-1 DO BEGIN
    temp= REFORM( tstack(*,*,ABS(matrices(i))) )
    IF ( matrices(i) LT 0 ) THEN temp= TRANSPOSE(temp)
    T= temp##T
ENDFOR

;
;Apply rotation
;
new_position= T ## new_position
rotation_matrix= T
;
;Maybe convert back from Cartesian into spherical
;
IF KEYWORD_SET(INTO_SPHERICAL) THEN BEGIN
    X= new_position(*,0)
    Y= new_position(*,1)
    Z= new_position(*,2)
    l= SQRT( X^2 + Y^2 )
    new_position(0,0)= SQRT( l^2 + Z^2 )
    new_position(0,1)= ATAN(z,l) * !radeg
    new_position(0,2)= ATAN(y,x) * !radeg ;Hapgood uses an ArcCos, but
this should work
ENDIF

RETURN,new_position
END

```

## PRO GEOSPACE\_CONVERT\_TEST

```

;Formats and units:
;   Day/Time format: YY/MM/DD HH.HHHHH
;   Degrees/Hemisphere format: Decimal degrees with 2 place(s).
;   Longitude 0 to 360, latitude -90 to 90.
;   Distance format: Earth radii with 5 place(s).
;
;polar
;   Time          GEI (RE)          GEO
(RE)             GM (RE)          GSE
(RE)             GSM (RE)         SM (RE)
;yy/mm/dd hh.hhhhh  X      Y      Z      X
Y      Z      X      Y      Z      X
Y      Z      X      Y      Z      X
Y      Z
;

```

```

;00/01/01 0.00000 1.89412 1.02594 6.89192 0.68262
-2.04311 6.89192 0.85108 0.00458 7.17039 -3.30372
2.49712 5.91512 -3.30372 0.82850 6.36693 -0.19481
0.82850 7.17039
;00/01/01 4.95000 1.85155 0.17499 -2.04050 -1.82577
-0.35419 -2.04050 0.14029 -1.84442 -2.04962 0.96501
1.70908 -1.94173 0.96501 1.83839 -1.81978 -0.20466
1.83839 -2.04962
;00/01/01 9.90000 7.25976 1.80121 3.82529 -4.29675
6.12262 3.82529 -7.74501 -2.15036 2.44379 -1.82681
7.70989 2.79316 -1.82681 7.99081 1.84148 -0.86959
7.99081 2.44379
;00/01/01 14.85000 5.14900 1.70083 7.19941 3.11398
4.43939 7.19941 -4.50089 4.35341 6.48277 -3.41217
5.86883 5.92878 -3.41217 5.95795 5.83921 -1.92691
5.95795 6.48277
;00/01/01 19.80000 -0.41009 0.40680 5.17012 -0.07490
0.57276 5.17012 -1.50754 0.10925 4.97787 -2.46367
0.04846 4.58167 -2.46367 -0.92680 4.48722 -1.19400
-0.92680 4.97787

```

```

;00/01/01 0.00000 1.89412 1.02594 6.89192 0.68262
-2.04311 6.89192 0.85108 0.00458 7.17039 -3.30372
2.49712 5.91512 -3.30372 0.82850 6.36693 -0.19481
0.82850 7.17039

```

```

CDF_EPOCH,epoch,2000,01,01,00,00,00,/COMPUTE

```

```

xgei= [1.89412 , 1.02594 , 6.89192 ]
xgeo= [0.68262 , -2.04311 , 6.89192 ]
xmag= [0.85108 , 0.00458 , 7.17039 ]
xgse= [-3.30372 , 2.49712 , 5.91512 ]
xgsm= [-3.30372 , 0.82850 , 6.36693 ]
xsm= [-0.19481 , 0.82850 , 7.17039 ]

```

```

PRINT,'MAG to GEO

```

```

',SQRT(TOTAL((xgeo-GEOSPACE_CONVERT(xmag,'mag','geo',CDF_EPOCH=epoch))^2))

```

```

PRINT,'GEO to GEI

```

```

',SQRT(TOTAL((xgei-GEOSPACE_CONVERT(xgeo,'geo','gei',CDF_EPOCH=epoch))^2))

```

```

PRINT,'GEI to GSE

```

```

',SQRT(TOTAL((xgse-GEOSPACE_CONVERT(xgei,'gei','gse',CDF_EPOCH=epoch))^2))

```

```

PRINT,'GSE to GSM

```

```

',SQRT(TOTAL((xgsm-GEOSPACE_CONVERT(xgse,'gse','gsm',CDF_EPOCH=epoch))^2))

```

```

PRINT,'GSM to SM

```

```

',SQRT(TOTAL((xsm-GEOSPACE_CONVERT(xgsm,'gsm','sm',CDF_EPOCH=epoch))^2))

```

```

PRINT,'SM to MAG

```

```

',SQRT(TOTAL((xmag-GEOSPACE_CONVERT(xsm,'sm','mag',CDF_EPOCH=epoch))^2))

```

```

PRINT,'GSM to GEO
',SQRT(TOTAL((xgeo-GEOSPACE_CONVERT(xgsm,'gsm','geo',CDF_EPOCH=epoch))^2))

PRINT,' '

;00/01/01 4.95000 1.85155 0.17499 -2.04050 -1.82577
-0.35419 -2.04050 0.14029 -1.84442 -2.04962 0.96501
1.70908 -1.94173 0.96501 1.83839 -1.81978 -0.20466
1.83839 -2.04962
CDF_EPOCH,epoch,2000,01,01,04,57,00,/COMPUTE
xgei= [ 1.85155 , 0.17499 , -2.04050]
xgeo= [-1.82577 , -0.35419 , -2.04050]
xmag= [0.14029 , -1.84442 , -2.04962 ]
xgse= [ 0.96501 , 1.70908 , -1.94173]
xgsm= [ 0.96501 , 1.83839 , -1.81978]
xsm= [-0.20466 , 1.83839 , -2.04962 ]

PRINT,'MAG to GEO
',SQRT(TOTAL((xgeo-GEOSPACE_CONVERT(xmag,'mag','geo',CDF_EPOCH=epoch))^2))
PRINT,'GEO to GEI
',SQRT(TOTAL((xgei-GEOSPACE_CONVERT(xgeo,'geo','gei',CDF_EPOCH=epoch))^2))
PRINT,'GEI to GSE
',SQRT(TOTAL((xgse-GEOSPACE_CONVERT(xgei,'gei','gse',CDF_EPOCH=epoch))^2))
PRINT,'GSE to GSM
',SQRT(TOTAL((xgsm-GEOSPACE_CONVERT(xgse,'gse','gsm',CDF_EPOCH=epoch))^2))
PRINT,'GSM to SM
',SQRT(TOTAL((xsm-GEOSPACE_CONVERT(xgsm,'gsm','sm',CDF_EPOCH=epoch))^2))

PRINT,'SM to MAG
',SQRT(TOTAL((xmag-GEOSPACE_CONVERT(xsm,'sm','mag',CDF_EPOCH=epoch))^2))
PRINT,'GSM to GEO
',SQRT(TOTAL((xgeo-GEOSPACE_CONVERT(xgsm,'gsm','geo',CDF_EPOCH=epoch))^2))

RETURN
END

```

---