Subject: Re: Satellite orbit procedure

Posted by Ed Wright on Thu, 28 Feb 2002 16:32:42 GMT

View Forum Message <> Reply to Message

in article 3C7E3189.88B08ED3@dlr.de, Richard Lord at Richard.Lord@dlr.de wrote on 2/28/02 5:32 AM:

> > Hi,

>

- > I'm looking for a procedure that takes as input the parameters of a
- > satellite orbit (Kepler orbit parameters), and outputs so-called
- > statevectors. A statevector contains the x,y,z and vx,vy,vz
- > coordinates at any particular point in time.

>

- > If anybody has seen a procedure that does something along the lines
- > explained above, I would be grateful to receive a copy.

>

- > Thanks in advance,
- > Richard

Give me another month and I'll finish testing a dlm interface to the JPL CSPICE library. The library includes routines for orbit computations, time conversion, and associated functions.

As always, Ed Wright JPL/NAIF

Subject: Re: Satellite orbit procedure
Posted by Craig Markwardt on Wed, 06 Mar 2002 14:50:41 GMT
View Forum Message <> Reply to Message

Richard Lord <Richard.Lord@dlr.de> writes:

> Hi,

>

- > I'm looking for a procedure that takes as input the parameters of a
- > satellite orbit (Kepler orbit parameters), and outputs so-called
- > statevectors. A statevector contains the x,y,z and vx,vy,vz
- > coordinates at any particular point in time.

>

- > If anybody has seen a procedure that does something along the lines
- > explained above, I would be grateful to receive a copy.

Hi Richard--

This kind of code can also be found in the HELIO procedure in the IDL

Astronomy Library. You have to be aware that this kind of code is only really good for bodies in the solar system. For near earth objects the story is a little different. The so-called "mean" orbital elements are also tied to an orbital model, like the SGP4. You would need to propagate the orbit using this model.

Subject: Re: Satellite orbit procedure
Posted by Liam E. Gumley on Wed, 06 Mar 2002 14:58:19 GMT
View Forum Message <> Reply to Message

Richard Lord wrote:

- > I'm looking for a procedure that takes as input the parameters of a
- > satellite orbit (Kepler orbit parameters), and outputs so-called
- > statevectors. A statevector contains the x,y,z and vx,vy,vz
- > coordinates at any particular point in time.

I looked into this problem a while ago. The most convenient solution I found was the "Satellite Tool Kit" (STK) software freely available from

http://www.stk.com/

The free version of their software accepts Keplerian elements (e.g. NORAD TLE format) and computes the corresponding state vectors propagated forward in time.

The STK software is unrelated to IDL.

Cheers, Liam. Practical IDL Programming http://www.gumley.com/