
Subject: Re: 2D non-linear, least-squares fitting routine needed

Posted by [rivers](#) on Fri, 25 Aug 1995 07:00:00 GMT

[View Forum Message](#) <> [Reply to Message](#)

In article <DDtt1x.uF@nntpa.cb.att.com>, windt@nanook.div111.att.com (David L. Windt) writes:

> I'm wondering if there are any IDL procedures available to perform

> 2D non-linear least-squares fitting, i.e., the equivalent of

> the CURVEFIT function (using a user-defined, multi-parameter

> FUNCTION) for 2-dimensions. It doesn't look like any of the

> fitting procedures contained in the standard library do that.

>

CURVEFIT can be used for 2D curve fitting. It just minimizes the sum of squares of residuals of a set of "observations" and "predictions". It doesn't really matter if those observations are on a 1-D, 2-D or n-D grid. You do need to call REFORM in the function which you write that CURVEFIT calls at each point, but that is trivial.

Mark Rivers	(312) 702-2279 (office)
CARS	(312) 702-9951 (secretary)
Univ. of Chicago	(312) 702-5454 (FAX)
5640 S. Ellis Ave.	(708) 922-0499 (home)
Chicago, IL 60637	rivers@cars3.uchicago.edu (Internet)
