
Subject: Re: Relative distances between vector elements and search for matches in other vector.

Posted by [David Fanning](#) on Wed, 25 Apr 2012 14:40:31 GMT

[View Forum Message](#) <> [Reply to Message](#)

Trifon Trifonov writes:

> I want to mention that I already made a working algorithm
> but I am 1000 % sure that the more experienced people
> here can help me to make it much more elegant and efficient.
> Later on I have a other problem on that I stuck but lets
> see can you help me on that and it is possible to solve
> it my self... if not I will post again :(

I have a reading assignment for you. :-)

Are FOR Loops the Embodiment of Pure Evil?

<http://www.idlcoyote.com/tips/forloops.html>

Are FOR Loops *Really* Evil?

<http://www.idlcoyote.com/tips/forloops2.html>

I don't have time to look at this in much detail,
except to point out that you have a lot of code
that looks like this:

```
for j=0L, n_elements(xx)-1 do begin
  ratio[j] = xx[0] - xx[j]
endfor
```

This code can *always* be written (and will work a LOT
faster) if you write it like this:

```
ratio = xx[0] - xx
```

Cheers,

David

--

David Fanning, Ph.D.

Fanning Software Consulting, Inc.

Coyote's Guide to IDL Programming: <http://www.idlcoyote.com/>

Sepore ma de ni thui. ("Perhaps thou speakest truth.")

Subject: Re: Relative distances between vector elements and search for matches in other vector.

Posted by [cgguido](#) on Wed, 25 Apr 2012 14:45:53 GMT

[View Forum Message](#) <> [Reply to Message](#)

What David said. Also, this line seems strange to me:

```
stel_dep = []
```

What are you trying to do there?

Gianguido

Subject: Re: Relative distances between vector elements and search for matches in other vector.

Posted by [cgguido](#) on Wed, 25 Apr 2012 15:03:35 GMT

[View Forum Message](#) <> [Reply to Message](#)

Also, I would also get `nx=n_elements(xx)` once and use it when I need it it.

Finally, your min for loop, I suspect one histogram could do it. but I am not sure what you're trying to do...

If you keep your loop then you could try this:

```
a = where( abs(ratio-ratio2[jjj]) LT 0.05 ,count)
```

and also avoid appending to `stel_dep` and `stel_wave` by creating them outside the loop and filling as needed. Then clip one at the end.

PS: histogram wise... I'd start with something approximately like:

```
nr=n_elements(ratio)
nr2=n_elements(ratio2)
```

```
bigratio=rebin(ratio,[nr,nr2],/sample)
bigratio2=transpose(rebin(ratio2,[nr2,nr], /sample))
```

now you have two arrays that are the same size, so you can do:

```
diff=bigratio-bigratio2
```

```
;good is now filled with 1s and 0s
good=abs(diff) LT 0.05
```

```
;you can count them by using DIMENSION with total:
count=total(good,2)
```

...

On Wednesday, April 25, 2012 9:03:00 AM UTC-5, Trifon Trifonov wrote:

> Hallo all
> I am new in IDL but I am increasing my level everyday and I am so happy that I discover IDL!!!
> I stuck very badly in one algorithm that I am developing for my Pipeline.
> I am searching for a routine or method to:
>
> 1st. define the relative distances between vector elements (e.g. distances between peaks in the spectrum(allready identified))
>
> 2nd. to identify the lines from another vector (bigger or smaller in array size)
> based on the criteria of the above (1st.)
>
> E.g.
>
> I have:
> `RATIO_STLINES` `FLOAT` = `Array[21]`
> 0.00000 -0.171038 -0.594733 -1.21922 -1.40923 -1.52994 -1.56168
> -1.92495 -2.28294 -2.34803 -2.38444
> -3.15909 -3.46812 -3.61399 -3.97221 -4.09896 -4.27114 -4.43637
> -5.69552 -6.05595 -6.23174
>
> `RATIO_CAT` `FLOAT` = `Array[44]`
> 0.00000 -0.209757 -0.249368 -0.930353 -1.05059 -1.21623 -1.21693
> -1.74758 -1.75778 -1.77888 -1.78749
> -2.03075 -2.22571 -2.99081 -3.29190 -3.31060 -3.42393 -3.43413
> -3.43844 -3.44364 -3.59758 -3.61648
> -3.76892 -3.80183 -3.92147 -4.26676 -4.53263 -5.02637 -5.31455
> -5.35194 -5.52650 -5.59132 -5.75987
> -5.81128 -6.60440 -6.68362 -6.71653 -6.73373 -6.81295 -6.95529
> -7.23017 -7.48834 -7.66608 -7.90535
>
> And the `RATIO_CAT` consist the lines from `RATIO_STLINES` (it is possible not all of them to be in the `RATIO_CAT`)
>
> So `RATIO_STLINES` is lets say shifted respect to `RATIO_CAT` and I want to identify lines based on their line ratios.
>
> I want to mention that I already made a working algorithm but I am 1000 % sure that the more experienced people here can help me to make it much more elegant and efficient. Later on I have a other problem on that I stuck but lets see can you help me on that and it is possible to solve it my self... if not I will post again :(

```

>
> Here is the code:
>
> ;Finding the peak of each line on the spectrum:
>
> peak = peaks(1-chip,0.9) ; finding the peaks from JJohnson IDL.
> ddd = xaxis[peak] ;wavelengths of the finded lines.
> lines = chip[peak] ;depts of the finded lines.
> ;xx - HITRAN lines
> ;z1- VALD catalog wavelengths
> ;z2 - VALD catalog line depths
>
> ;Finding the relative distance between the HITRAN lines (nm):
> ratio=fltarr(n_elements(xx))
>
> for j=0L, n_elements(xx)-1 do begin
>   ratio[j] = xx[0] - xx[j]
>   endfor
>
> ;Finding the relative distance between each line on the spectrum:
> ratio2=fltarr(n_elements(ddd))
>
> for jj=0L, n_elements(ddd)-1 do begin
>   ratio2[jj] = ddd[0] - ddd[jj]
>   endfor
>
> ;Finding the relative distance between the lines from the VALD catalog:
> ratio_vald=fltarr(n_elements(z1))
>
> for jjj=0L, n_elements(z1)-1 do begin
>   ratio_vald[jjj] = z1[0] - z1[jjj]
>   endfor
>
>
> ; Identify the telluric lines on the spectra. Only stellar lines will be left.
> telluric_depts = fltarr(n_elements(xx))
> telluric_waves = fltarr(n_elements(xx))
> stel_dep = []
> stel_wave = []
> stel_ra = []
> for jjj = 0L, n_elements(ratio2)-1 do begin
>   a = where((ratio GT (ratio2[jjj] - 0.05)) and (ratio LT (ratio2[jjj] + 0.05)),count)
>
>   if count gt 0d0 then begin
>     telluric_depts[a] = lines[jjj]
>     telluric_waves[a] = ddd[jjj]
>   endif
>

```

```

> if count le 0d0 then begin
>   stel_dep =[stel_dep, lines[jjj]]
>   stel_wave =[stel_wave, ddd[jjj]]
>   ;stel_ra = [stel_ra, ratio2[jjj]]
>   endif
>
> endfor
>
> ;Finding the relative distance between the lines that left from the science spectrum:
>
> ratio_stlines = fltarr(n_elements(stel_wave))
> for k = 0L, n_elements(stel_wave) - 1 do begin
>   ratio_stlines[k] = stel_wave[0] - stel_wave[k]
> endfor
>
>
> And I will stop here because I have more problems below...
>
> Thank you very much in advance!

```

Subject: Re: Relative distances between vector elements and search for matches in other vector.

Posted by [Tito](#) on Thu, 26 Apr 2012 13:48:09 GMT

[View Forum Message](#) <> [Reply to Message](#)

First of all I want to thank you for you input!

@ David: ratio = xx[0] - xx ... How it is possible to be sooo easy?!?
I corrected this and now looks much better and understandable.

@ Gianguido: a = where(abs(ratio-ratio2[jjj]) LT 0.05 ,count) - this works the same as mine but it is much shorter! thanks for that!

I tested: good=abs(diff) LT 0.05 and count=total(good,2),

And this gives me, indexes of 1 and 0, and I want the indexes of ratio2, like:

0,1,2,4,5,8,9 and i get :

print,good

0,19,38,....719

The loop is working however thats maybe because the first line of the catalog and the first line of the spectrum "this time" are the same. e.g.

ratio = xx[0] - xx

0,-0,17,-0,34.....

and

ratio2 = ddd[0] - ddd

0,-0,1703,-0,22,-0,3403.... so it is easy to identify those differences which pass the (a = where(

abs(ratio-ratio2[jjj]) LT 0.05 ,count)) criteria.

in other hand if they are shifted e.g. the first line from catalog is not the first identified line in the spectrum I am having a problem...

My goal is to identify lines from different catalogs with those from my spectrum based on their relative differences(distances) this is necessary because usually the lines from the spectrum are shifted but they have the same ratios between each other (they might differ slightly) with those from the catalog.

So first I am finding the atmosphere lines on the spectra and what is left is the stellar lines which I will search again in catalog.

This is important for me because

1st. it will be automatic! ,

2nd. the method will be more accurate then if I have to select the lines manually. Sometimes there is 2 or more lines very close to each other and selecting is pain.

Now i probably need something that will find the same space between the two vectors.

I read about histogram from David's website but my head is about to explode.

I am still investigating, how to use 'histogram'...

Lets assume that I identify the telluric lines and now I have only the stellar which I will search from catalog and this time I know that the first line of the spectrum and the catalog are not the same.

So I did this:

```
.....
selected_dep = [] ; create empty arrays because I want to append (I dont know the result number
of array elements)
selected_lines = []
ratio_sel_vald = []
for z = 0L,n_elements(ratio_stlines)-1 do begin
  for zz = 0L,n_elements(ratio_stlines)-1 do begin

    aa = where( abs((ratio_stlines[zz] - ratio_stlines[z]) - ratio_vald) LT 0.0025, count)

    if count gt 0d0 then begin
      selected_dep = [selected_dep,z2[aa]]
      selected_lines = [selected_lines, z1[aa]]
      ratio_sel_vald = [ratio_sel_vald,ratio_vald[aa]]

    endif
  endfor
endfor
```

;;;I have to use 'Uniq' because from this 2 loops I have many Duplicates.

```
selected_lines = selected_lines[uniq(selected_lines,sort(selected_lines))]
```

```
ratio_sel_vald = ratio_sel_vald[uniq(ratio_sel_vald,sort(abs(ratio_sel_vald)) )]
```

And this is working! if I want to find which lines from the catalog are in my spectra. BUT i want to give to "selected_lines" not the "selected_dep"(depths) which are from the catalog, I want to associate the fonded lines with the depths from the spectra. (- And this is my biggest problem actually I dont know how to do this!)

Later I have to construct synthetic template(spectrum) based on that data, that will have the same depths and profile like those from the spectra BUT different wavelength. Then I will cross-correlate both synthetic mask + spectra in order to find the shift.

Thank you again for your time reading this and trying to help me!

Trifon

Subject: Re: Relative distances between vector elements and search for matches in other vector.

Posted by [cgguido](#) on Thu, 26 Apr 2012 21:30:44 GMT

[View Forum Message](#) <> [Reply to Message](#)

Trifon, you're most welcome.

If you can estimate the max possible size of your result (even if you overestimate it by a lot) I would do:

```
selected_dep=fltarr(1e4)*0-1
;this creates a huge array with "impossible values" -1.
Then you fill it as you go with real values, And at the end just do:
selected_dep = selected_dep[ where(selected_dep NE -1)]
This is usually much faster!
```

In general, I am still confused about what you are trying to do. Could you give a simple example of input and output? Something like:

```
;inputs:
r1=[1,3,3,4,5]
r2=[2,-1,4,3]
```

```
;outputs:
3,4
```

Don't bother with technical (geological) terms, they just confuse non-specialists. Just give us the simple examples and we'll see if we can speed this up for you a bit.

Gianguido
