
Subject: Re: locating the closest values?

Posted by on Tue, 26 Jun 2012 21:28:56 GMT

[View Forum Message](#) <> [Reply to Message](#)

Den tisdagen den 26:e juni 2012 kl. 22:55:34 UTC+2 skrev anil:

> Hi,
> I have two datasets/files. 1st one contains the values i want to find
> (latitudes and longitudes) and the second file contains 3 columns as
> longitude latitude and depth.
> What i want to do is to read my desired values from file 1 and find
> the closest values in file 2 and the corresponding 3rd column(depth)
> from file 2. here is what i did:
> 1st file:
> 42.25 30.57
> 41.17 32.58
>
> 2nd file
> 42.55 34.42 1000
> 41.54 33.21 1500
>
> what i did is :
>
> difference1=abs(lon-desiredlon)
> a1=min(difference1,index1)
>
> difference2=abs(lat-desiredlat)
> a2=min(difference2,index2)
>
> in this case, lon(index1) and lat(index2) become my desired values. no
> problem with that. but how do i get the 3rd column. i could not match
> index1 and index2 with where function.
> what am i missing here?

Index1 and index2 are different because being close in one of lat and lon is not the same as being closest in both. I suspect what you want is something like
 $a = \min(\sqrt{\text{difference1}^2 + \text{difference2}^2}, \text{index})$, which should give you the index to the point with the shortest distance from the desired point. (You may want to scale the differences to length units first.)

Subject: Re: locating the closest values?

Posted by [anil](#) on Tue, 26 Jun 2012 21:50:29 GMT

[View Forum Message](#) <> [Reply to Message](#)

On Jun 27, 12:28 am, Mats Löfdahl <mats.lofd...@gmail.com> wrote:

> Den tisdagen den 26:e juni 2012 kl. 22:55:34 UTC+2 skrev anil:
>
>

```

>
>
>
>
>
>
>
>> Hi,
>> I have two datasets/files. 1st one contains the values i want to find
>> (latitudes and longitudes) and the second file contains 3 columns as
>> longitude latitude and depth.
>> What i want to do is to read my desired values from file 1 and find
>> the closest values in file 2 and the corresponding 3rd column(depth)
>> from file 2. here is what i did:
>> 1st file:
>> 42.25 30.57
>> 41.17 32.58
>> .....
>> 2nd file
>> 42.55 34.42 1000
>> 41.54 33.21 1500
>> .....
>> what i did is :
>
>> difference1=abs(lon-desiredlon)
>> a1=min(difference1,index1)
>
>> difference2=abs(lat-desiredlat)
>> a2=min(difference2,index2)
>
>> in this case, lon(index1) and lat(index2) become my desired values. no
>> problem with that. but how do i get the 3rd column. i could not match
>> index1 and index2 with where function.
>> what am i missing here?
>
> Index1 and index2 are different because being close in one of lat and lon is not the same as
> being closest in both. I suspect what you want is something like
> a=min(sqrt(difference1^2+difference2^2),index), which should give you the index to the point with
> the shortest distance from the desired point. (You may want to scale the differences to length
> units first.)

```

Thank you for your answer. I guess I just found it. it is as simple as it is: where function :) :

c=where(lon eq lon(index1) and lat eq lat(index2))
 this c is what i want. just the statement giving the exact locations.

Subject: Re: locating the closest values?

Posted by [cgguido](#) on Tue, 26 Jun 2012 22:25:24 GMT

[View Forum Message](#) <> [Reply to Message](#)

I'd use `measure_distance` to get them all in one go.... Then search for the minima...

Subject: Re: locating the closest values?

Posted by [Russell Ryan](#) on Wed, 27 Jun 2012 01:35:03 GMT

[View Forum Message](#) <> [Reply to Message](#)

On Tuesday, June 26, 2012 5:28:56 PM UTC-4, Mats Löfdahl wrote:

> Den tisdagen den 26:e juni 2012 kl. 22:55:34 UTC+2 skrev anil:

>> Hi,

>> I have two datasets/files. 1st one contains the values i want to find
>> (latitudes and longitudes) and the second file contains 3 columns as
>> longitude latitude and depth.

>> What i want to do is to read my desired values from file 1 and find
>> the closest values in file 2 and the corresponding 3rd column(depth)
>> from file 2. here is what i did:

>> 1st file:

>> 42.25 30.57

>> 41.17 32.58

>>

>> 2nd file

>> 42.55 34.42 1000

>> 41.54 33.21 1500

>>

>> what i did is :

>>

>> `difference1=abs(lon-desiredlon)`

>> `a1=min(difference1,index1)`

>>

>> `difference2=abs(lat-desiredlat)`

>> `a2=min(difference2,index2)`

>>

>> in this case, `lon(index1)` and `lat(index2)` become my desired values. no

>> problem with that. but how do i get the 3rd column. i could not match

>> `index1` and `index2` with `where` function.

>> what am i missing here?

>

> `Index1` and `index2` are different because being close in one of lat and lon is not the same as being closest in both. I suspect what you want is something like

`a=min(sqrt(difference1^2+difference2^2),index)`, which should give you the index to the point with the shortest distance from the desired point. (You may want to scale the differences to length units first.)

So, that's going to work, but the distance you get "a" is probably not the minimum distance since

you're working with angles. That said, you should look up the relevant equation for "great circle" distances to get the true minimum.

Subject: Re: locating the closest values?

Posted by [cgguido](#) on Wed, 27 Jun 2012 15:51:19 GMT

[View Forum Message](#) <> [Reply to Message](#)

Ah yes, pesky distances on a sphere.... Also, I meant to say distance_measure.

G
