
Subject: Re: regridding questions...

Posted by [safier](#) on Fri, 19 Apr 1996 07:00:00 GMT

[View Forum Message](#) <> [Reply to Message](#)

>>>> > "Phil" == Phil Williams <williams@irc.chmcc.org> writes:

Phil> I have some data that is sampled on a close-to-rectangular
Phil> grid that looks something like the following:

Phil> _____ / /\ \ _____ \ \ / _____ / /\ /

Phil> Are you following me? I have the x and y points defined in
Phil> an MxN array. The data is an MxN complex array. How would
Phil> I go about using, or can I even, some IDL function to
Phil> accomplish getting the data points on a regular rectangular
Phil> grid?

Phil> I have tried to use triangulate and trigrd with no success.
Phil> krig2d can't allocate enough space for the new array.

Phil> I've come to a dead end and any help would be greatly
Phil> appreciated!

Phil> TIA,

How about using bicubic spline interpolation to interpolate the values
on a nice rectangular grid of your choice?

Cheers,

Pedro

--

Pedro N. Safier | "God offers to everyone his
Department of Astronomy | choice between truth and repose.
U. of Maryland at College Park | Take which you please--you can
phone: 301-405-1531; fax: 301-314-9067 | never have both." R. W. Emerson
