
Subject: Re: a_correlate is wrong...

Posted by [B}rd Krane](#) on Thu, 11 Dec 1997 08:00:00 GMT

[View Forum Message](#) <> [Reply to Message](#)

William Connolley wrote:

>

> The a_correlate function seems to be wrong to me. Whats wrong is that
> at every lag, the correlation is divided by the total number of elements
> in the series, not by the number-in-series minus the lag. If you take
> a perfectly correlated series, this make the correlations at higher lags
> get artificially smaller.

This is not necessarily wrong, it just gives you a biased estimator
instead of an unbiased estimator. The autocovariance (correlation) is
defined as the ensemble average of two elements separated by k elements

$$R[k] = \langle x[i+k] x[i] \rangle$$

With a finite record length the average for the maximum lag contains
only one entry and is highly unreliable. Thus it is sometimes given
a smaller weight, i.e. divide by N instead of 1.

This is a sloppy explanation, but it should give you the necessary
starting points for further reading, e.g.

Modern Spectral Estimation

Steven M. Kay

ISBN 0-13-598582-x

Baard

B{aa}rd Krane | Email : bard.krane@fys.uio.no
Institute of Physics | Phone : (+47) 2285 5666
University of Oslo | Fax : (+47) 2285 5671
POB 1048, blindern, N-0316 Norway